

THE MERCK INDEX

TENTH EDITION

THE MERCK INDEX

AN ENCYCLOPEDIA OF
CHEMICALS, DRUGS, AND BIOLOGICALS

TENTH EDITION

Martha Windholz, *Editor*
Susan Budavari, *Co-Editor*
Rosemary F. Blumetti, *Associate Editor*
Elizabeth S. Otterbein, *Assistant Editor*

0002

Published by
MERCK & CO., INC.
RAHWAY, N. J., U. S. A.

1983

Copyright © 1983 by Merck & Co., Inc.

Previous Editions

Copyright © 1940, 1952, 1960, 1968, 1976
by Merck & Co., Inc.

All rights reserved. Copyright under the Universal Copyright
Convention and the International Copyright Convention.
Copyright reserved under the Pan-American Copyright
Convention.

Merck & Co., Inc.

Rahway, New Jersey, U.S.A.

MERCK SHARP & DOHME
West Point, Pa.

MERCK SHARP & DOHME INTERNATIONAL
Rahway, N. J.

MERCK SHARP & DOHME RESEARCH LABORATORIES
Rahway, N. J. / West Point, Pa.

MSD AGVET DIVISION
Rahway, N. J.

HUBBARD FARMS, INC.
Walpole, N. H.

MERCK CHEMICAL MANUFACTURING DIVISION
Rahway, N. J.

MERCK CHEMICAL DIVISION
Rahway, N. J.

KELCO/AIL INTERNATIONAL LIMITED
London, England

BALTIMORE AIRCOIL COMPANY, INC.
Baltimore, Md.

CALGON CARBON CORPORATION
Pittsburgh, Pa.

KELCO DIVISION
San Diego, Calif.

7000


1st Edition – 1889
2nd Edition – 1896
3rd Edition – 1907
4th Edition – 1930
5th Edition – 1940
6th Edition – 1952
7th Edition – 1960
8th Edition – 1968
9th Edition – 1976
10th Edition – 1983

Library of Congress Catalog
Card Number 83-61075
ISBN Number 911910-27-1

Printed in the U.S.A.


First Printing – October 1983
Second Printing – March 1984

Asta Chemische Fabrik); cf. Goldberg, Shapero, *J. Pharm. Pharmacol.* 6, 171 (1954).



Free base. Insol in water. Hydrochloride, $C_{23}H_{27}NO_3 \cdot HCl$, crystals, mp 199-200°. Sparingly sol in water (1:500). Aq solns are acid to litmus. THERAP CAT: Antispasmodic.


6597. Oethilnone. 2-Octyl-3(2H)-isothiazolone; 2-octyl-4-isothiazolin-3-one; RH-893; Kathon. $C_{11}H_{19}NOS$; mol wt 213.34. C 61.93%, H 8.97%, N 6.57%, O 7.50%, S 15.03%. Prepn: S. N. Lewis *et al.*, Fr. pat. 1,555,416 corresp to U.S. pat. 3,761,488 (1969, 1973 to Rohm & Haas); *idem*, *J. Heterocycl. Chem.* 8, 571 (1971).



Liquid, bp_{0.01} 120°. uv max (methanol): 280 nm (log ϵ 3.88).

USE: Fungicide. Biocide in cooling-tower water, paints, cutting oils, cosmetics and shampoo; for leather preservation.

6598. Octodrine. 6-Methyl-2-heptanamine; 6-methyl-2-heptylamine; 2-methyl-6-aminoheptane; 6-amino-2-methylheptane; 2-amino-6-methylheptane; α,ϵ -dimethylhexylamine; 1,5-dimethylhexylamine; SKF 51; Vaporpac. $C_8H_{19}N$; mol wt 129.24. C 74.34%, H 14.82%, N 10.84%. Prepd from the corresponding saturated ketone: Rohrmann, Shonle, *J. Am. Chem. Soc.* 66, 1516 (1944). Pharmacology: E. J. Fellows, *J. Pharmacol. Exp. Ther.* 90, 351 (1947).




dl-Form, viscous liquid, fishy odor, bp 154-156°. n_D^{20} 1.4200.

Hydrochloride, $C_8H_{19}N \cdot HCl$, crystals, sol in water. LD₅₀ i.p. in mice, rats: 59, 41.5 mg/kg. E. J. Fellows, *loc. cit.*

Sulfate, $2C_8H_{19}N \cdot H_2SO_4$, crystals, sol in water. THERAP CAT: Adrenergic (vasoconstrictor), local anesthetic.

6599. Octopamine. α -(Aminomethyl)-4-hydroxybenzene-methanol; α -(aminomethyl)-*p*-hydroxybenzyl alcohol; 1-(*p*-hydroxyphenyl)-2-aminoethanol; norysympatol; norysinephrine; *p*-hydroxyphenylethanolamine; WV 569. $C_9H_{11}NO_2$; mol wt 153.18. C 62.72%, H 7.24%, N 9.14%, O 20.89%. Biogenic amine formed by β -hydroxylation of tyramine by the enzyme dopamine β -hydroxylase: Pisano *et al.*, *Biochim. Biophys. Acta* 43, 566 (1960). Identification: Erspamer, *Nature* 169, 375 (1952). Found in the salivary glands of *Octopus vulgaris*, *O. macropus*, and of *Eledone moschata*: *idem*, *Arzneimittel-Forsch.* 2, 253 (1952); in mammalian nerves: Molinoff, Axelrod, *Science* 164, 428 (1969); in cockroach nervous system: Nathanson, Greengard, *ibid.* 180, 308 (1973). Prepd synthetically: Asscher, U.S. pat. 2,585,988 (1952). The natural *D*(-)-form is 3 times more potent than the *L*(+)-form in producing cardiovascular adrenergic responses in anesthetized dogs and cats. Korol, Soffer, *The Pharmacologist* 5, 247 (1963). Prepn of *D*- and *L*-forms: Kappe, Armstrong, *J. Med. Chem.* 7, 569 (1964). In invertebrate nervous systems octopamine may function as a neurotransmitter: Saavedra *et al.*, *Science* 185, 364 (1974).




D(-)-Form, crystals from hot water which change at about 160° to a compd which melts at >250° (dec). $[\alpha]_D^{25}$ -56.0° (0.1 *N* HCl); -37.4° (H_2O).

DL-Form hydrochloride, $C_8H_{12}ClNO_2$, *Epiprenor*, *Norden*, *Norfen*, *Norphen* (ampules). Crystals, dec 170°. Freely sol in water.

THERAP CAT: Adrenergic.


6600. Octotiamine. 6-(Acetylthio)-8-[[2-[(4-amino-2-methyl-5-pyrimidinyl)methyl]formylamino]-1-(2-hydroxyethyl)-1-propenyl]dithiooctanoic acid methyl ester; 8-[[2-[[N-[(4-amino-2-methyl-5-pyrimidinyl)methyl]formamido]-1-(2-hydroxyethyl)propenyl]dithio]-6-mercaptooctanoic acid methyl ester (Stor 6)-acetate; S-(3-acetylthio-7-carbomethoxyheptylthio)thiamine; thiamine 8-(methyl 6-acetyldihydrothioate) disulfide; Gerostop; Neuvitan; TATD. $C_{29}H_{36}N_4O_6S_3$; mol wt 544.87. C 50.72%, H 6.66%, N 10.28%, O 14.68%, S 17.66%. Prepn: Ohara *et al.*, U.S. pat. 3,098,856 (1963 to Fujisawa).



Crystals, mp 106-109°. uv max: 234, 277 nm (ϵ 16,200, 5820).

Hydrochloride, $C_{29}H_{36}N_4O_6S_3 \cdot HCl$, crystals from ether + abs ethanol, mp 134.5-135°. uv max: 233 nm (ϵ 23,000). THERAP CAT: Long-acting oral thiamine.

6601. Octoxynol. α -[4-(1,1,3,3-Tetramethylbutyl)phenyl]- ω -hydroxypoly(oxy-1,2-ethanediyl); octylphenoxy polyethoxyethanol; polyethylene glycol *p*-isooctylphenyl ether. Prepd by reacting isooctylphenol with ethylene oxide. Refs, nomenclature, see nonoxynol. Trademarks for series of octoxynols include *Igepal CA*, *Polytergent G*, *Triton X*.



Octoxynol (N.F.), mixture in which *n* ranges from 5 to 15; average comp. (*n* = 10): $C_{34}H_{62}O_{12}$; av. mol wt 647. Pale yellow, viscous liquid. d_4^{25} 1.0595. n_D^{25} 1.4894. Miscible with water, alcohol, acetone. Sol in benzene, toluene. Insol in petr ether. pH of 5% aq soln: 7-9. *Octoxynol-9* (USAN), average comp. (*n* = 9): $C_{32}H_{58}O_{11}$. Trademarks of products where *n* = 9 to 10: *Conco NIX-100*, *Igepal CA-630*, *Neutro-nyx 605*, *Triton X-100*. Ingredient of *Preceptin*.

USE: Nonionic detergent, emulsifier, dispersing agent. Ingredient of nitrofurazone soln, N.F.

THERAP CAT: Spermatocide.

6602. Octyl Acetate. Acetic acid α -ethylhexyl ester; 2-ethylhexyl acetate. $C_{10}H_{20}O_2$; mol wt 172.26. C 69.72%, H 11.70%, O 18.58%. $CH_3(CH_2)_3CH(C_2H_5)CH_2OOCCH_3$.

Liquid. d_4^{20} 0.873. bp 199°. Solidifies about -80°. n_D^{20} 1.4204. Flash pt, open cup: 190°F (88°C); closed cup: 56°F (13°C). Very slightly sol in water; misc with alcohol, oils, and other organic liquids. LD₅₀ orally in rats: 3.0 g/kg, H. F. Smyth, C. P. Carpenter, *J. Ind. Hyg. Toxicol.* 26, 269 (1944).

USE: Solvent for nitrocellulose, some resins, waxes, and oils.

6603. *n*-Octyl Bromide. 1-Bromooctane. $C_8H_{17}Br$; mol wt 193.13. C 49.75%, H 8.87%, Br 41.38%. $CH_3(CH_2)_6CH_2Br$. Prepd from hydrobromic acid and *n*-octanol: O.